

Estimation of carbon balance components over agricultural landscapes based on remote sensing and field measurements

Thesis of PhD Dissertation

Györgyi Gelybó

Eötvös Loránd University
Faculty of Science

Doctoral School of Earth Sciences
Head of the doctoral school: *Dr. József Nemes-Nagy, professor*

Geography-Meteorology Program
Head of the PhD Program: *Dr. Mária Szabó, professor*



Advisors:

Dr. Judit Bartholy

Dr. Zoltán Barcza

Department of Meteorology
Institute of Geography and Earth Sciences, Eötvös Loránd University

Budapest, 2014

Introduction and objectives

Biogeochemical cycles in general and the biosphere-atmosphere carbon exchange in particular are of current high priority research areas for the understanding of climate change. Uncertainties in future climate scenarios need a more accurate incorporation of biosphere-atmosphere interactions because of the two-way interaction between the carbon cycle and the climate system. Oceans and the terrestrial ecosystems currently act as a net sink of atmospheric CO₂ partly compensating for anthropogenic emissions. One of the most important scientific questions is whether this sink capacity can be maintained or even increased under changing environmental conditions.

Remote sensing applications provide spatially explicit information on land surface (including vegetation state and activity), representing a valuable source of measurement based data for carbon cycle modeling. In my PhD research I am dealing with the possible applications of satellite remote sensing in estimating vegetation-atmosphere CO₂ exchange using a data-oriented model focusing especially on agroecosystems. Based on remote sensing data it is possible to estimate the gross carbon uptake of plants (Gross Primary Production; GPP) which is one of the most important and highly uncertain carbon cycle components of terrestrial ecosystems. The two main objectives of the dissertation are (i) to investigate the accuracy of remotely sensed GPP estimations against *in situ* measurements carried out over Hungarian and international agricultural fields and (ii) to improve model accuracy based on model parameter optimization.

Data and methods

The MOD17A2 product that is built upon data provided by the MODIS¹ sensor onboard NASA's² *Terra* and *Aqua* satellites provides *Net Photosynthesis and GPP* data since 2000 with 1 km spatial, and 8 days temporal resolution. The underlying MOD17 model is based on the LUE³ approach and requires weather data to simulate environmental stress on plant functioning. The 5-parameter model considers minimum temperature (T_{\min}) and atmospheric vapour pressure deficit (VPD) data to reduce maximum LUE (ϵ_{\max}) of the plant according to the actual environmental circumstances. GPP data is calculated in the model based on other MODIS data products that describe land cover information and vegetation activity.

Evaluation of remote sensing GPP estimations is usually carried out using field measurements of GPP as validation and calibration dataset. Application of GPP data from eddy covariance (EC) measurements is a common and frequently applied way to assess gross carbon uptake based on field measurements on the plot level. The actual representativeness of the EC measurements is influenced by several factors, such as measurement height (h), wind direction, atmospheric stability, orography, etc. The so-called footprint, or source area of the EC tower gives the area from which the measured CO₂ flux originates. Its location varies with time depending on atmospheric circumstances, but when integrated over a longer time period it provides information

¹ MODerate Resolution Imaging Sctroradiometer

² National Aeronautics and Space Administration

³ Light Use Efficiency

on the contribution of the neighboring areas to the measured flux. In general, the extent of the footprint is greater in case of a taller tower (relative to the canopy height).

In my PhD work, I used two types of EC data: data from a tall tower in Hungary and data from selected members of a network of short-tower based EC sites (FLUXNET) across Europe and the US.

In Hungary (near Hegyhátsál, Vas county) continuous measurements of carbon dioxide fluxes with eddy-covariance method have been taking place since 1997 at 82 m height (FLUXNET site ID: HU-He1). Being a tall tower, HU-He1 provides regional scale information on the surrounding heterogeneous agricultural landscape, hence it is expected that representativeness of the data is closer to the 1 km size of the MODIS pixel than in case of a shorter tower. 6 measurement years were used in the study to explore the effect of landscape heterogeneity on validation.

For calibration of MOD17 short-tower ($h < 30$ m above the canopy) based EC tower sites have been selected over croplands providing nearly 60 site-years of GPP data. C3 and C4 type cereals are grown on the study sites. After standardized data processing, I used this dataset to perform multi-site validation and calibration of the MOD17 model.

Results

The main results of the PhD work are as follows:

1. A coherent and standardized database of GPP data from agricultural EC validation sites was built. Four different methods were implemented to fill data gaps in EC measurements and to retrieve GPP data. Based on the four estimates of GPP, uncertainty of GPP data was assessed. The database also contains ancillary data such as meteorological measurements and crop information in standardized format. The database is also suitable for future validation/calibration of other ecosystem models.
2. Adaptation of the MOD17 model was carried out to be able to modify model setup and model parameters.
3. The first validation of the MOD17 model for Hungarian agricultural landscape has been performed, also considering the special properties of the site. The validation included four steps in four model setups: (i) the original model setup (ii) setup to account for errors of the input data (iii) setup for the rough consideration of tower footprint over the heterogeneous landscape, (iv) downscaling of the model using 250 m resolution NDVI data combined with footprint information. The results suggested underestimation of cropland validation data by original MOD17 estimates.
4. It was shown that, patchiness of the vegetation in the footprint considerably affects validation results, which means that mismatch of spatial representativeness of remote sensing and *in situ* data need to be accounted for. However, the dynamic consideration of temporal

changes of the footprint combined with a downscaling of the model to the approximate spatial scale of the individual crop fields in the region improved model-measurement agreement. Due to downscaling modeling efficiency (ME) increased from 0.783 to 0.884, root mean square error (RMSE) decreased from $1.095 \text{ gC m}^{-2} \text{ day}^{-1}$ to $0.815 \text{ gC m}^{-2} \text{ day}^{-1}$, although GPP remained underestimated (bias decreased from $-0.680 \text{ gC m}^{-2} \text{ day}^{-1}$ to $-0.426 \text{ gC m}^{-2} \text{ day}^{-1}$). This suggests that spatial mismatch is not solely responsible for the disagreement.

5. A multi-site validation of the MOD17 model has been carried out over croplands. The results showed a significant underestimation of measured (EC) data. The underestimation was more pronounced in case of C4 than C3 plant types. This raises the need for a crop specific separation of the current single general crop type in the MOD17 model. To our knowledge, such an extensive, multi-site evaluation of the model over agricultural vegetation has not been reported in the scientific literature so far.
6. Sensitivity analysis and model calibration of MOD17 was carried out using the GLUE⁴ method, which is a Monte-Carlo based global optimization approach. The model turned out to be most sensitive to the ϵ_{\max} parameter while other parameters (describing T_{\min} and VPD stress) barely influence model result.
7. The optimized parameter sets, obtained via a crop specific calibration of the MOD17 model for C4 and C3 plants separately, resulted in

⁴ Generalized Likelihood Uncertainty Estimation

higher ϵ_{\max} for C4 than for C3 crops ($0.002527 \text{ kg C MJ}^{-1}$ and $0.001487 \text{ kg C MJ}^{-1}$, respectively). The validation of the optimized MOD17 model for C4 and C3 plants separately resulted in a much better agreement between remote sensing and EC tower based estimates ($R^2 = 0.356$ ($p=0.001$) and 0.71 ($p \leq 10^{-7}$)).

Conclusions

The first validation results of the MOD17 model for Hungarian croplands revealed that an appropriate methodology is required in order to obtain defensible validation results at regional scale. The results demonstrate that careful application of remote sensing based information can provide reasonable estimate on cropland GPP for both C3 and C4 plants, especially if consideration of differences between – at least – basic physiological properties of crops is introduced in the model. Due to the multi-site approach, the results can contribute to a more effective use of the MOD17 remote sensing product for croplands over large regions of the USA and Europe. Croplands occupy about 45 % of the European and 18 % (35% together with pastures and rangeland) of the USA land surface, clearly reflecting their significance in the continental carbon balance.

In Hungary, availability of reliable data on the spatial distribution of C3 and C4 crops, however, can limit applicability of the crop-specific approach described above. Therefore, data collection and the application of remote sensing data at the appropriate spatial scale is an important and challenging task in the future to support Hungarian carbon balance estimations.

Publications related to this thesis

Papers

- Gelybó Gy., Barcza Z., Kern A., Kljun N. (2013) Effect of spatial heterogeneity on the validation of remote sensing based GPP estimations. *AGRICULTURAL AND FOREST METEOROLOGY* 174–175: 43–53. doi:10.1016/j.agrformet.2013.02.003 (IF: 3.421 (2012))
- Tóth E., Gelybó Gy., Kása I., Farkas Cs. (2013) A művelés hatása a talaj szén-dioxid kibocsátására: II. A talaj vízpotenciál értéke és CO₂ emissziója közötti összefüggések vizsgálata laboratóriumi módszertani tesztelés során. *AGROKÉMIA ÉS TALAJTAN* 62:(2) pp. 299–310.
- Barcza Z., Haszpra L., Hidy D., Gelybó Gy., Dobor L. (2012) A légkör és a bioszféra közötti szén-dioxid csere vizsgálata. *LÉGKÖR* 57:(4) pp. 147–149.
- Mihon D., Bacu V., Gorgan D., Mészáros R., Gelybó Gy. (2010) Practical considerations on the GreenView application development and execution over SEE-GRID. *EARTH SCIENCE INFORMATICS* 3:(4) pp. 247–258. doi:10.1007/s12145-010-0066-z (IF:0.657)

Book chapters

- Tóth E., Gelybó Gy., Farkas Cs. (2013) Talajművelés hatása a mezőgazdasági talajok szén-dioxid kibocsátására. In: Janda T (Ed.) II. ATK Tudományos Nap: Velünk Élő Tudomány (A Magyar Tudomány Ünnepe alkalmából) Konferencia helye, ideje: Martonvásár, Magyarország, 2013.11.08 Martonvásár: MTA Agrártudományi Kutatóközpont, 2013. pp. 144–151. (ISBN: 978-963-8351-41-8)
- Gelybó Gy., Tóth E., Barcza Z., Farkas Cs., Kern A. (2011) Carbon cycle of agricultural lands in Hungary from patch to regional scale. In: A Celkova (Ed.) 19th International Poster Day Transport of Water, Chemicals and Energy in the Soil-Plant-Atmosphere System. Bratislava, Szlovákia, 2011.11.09 Bratislava: Institute of Art History of SAS, 2011. pp. 115–126. (ISBN: 978-80-89139-26-2)
- Tóth E., Barcza Z., Birkás M., Gelybó Gy., Zsembeli J., Bottlik L., Davis J. D., Haszpra L., Kern A., Kljun N., Koós S., Kovács Gy., Stingli A., Farkas Cs. (2010) Measurements and estimations of biosphere-atmosphere exchange of greenhouse gases: Arable lands. In: Haszpra L. (Ed.) *Atmospheric Greenhouse Gases: The Hungarian Perspective*. Dordrecht; London; New York: Springer-Science, 2011. pp. 157–197. (ISBN:978-90-481-9949-5)
- Somogyi Z., Hidy D., Gelybó Gy., Barcza Z., Churkina G., Haszpra L., Horváth L., Machon A., Grosz B. (2010) Modeling of biosphere-atmosphere

- exchange of greenhouse gases: Models and their adaptation In: Haszpra L. (Ed.) Atmospheric Greenhouse Gases: The Hungarian Perspective. Dordrecht; London; New York: Springer-Science, 2011. pp. 201-208. (ISBN:978-90-481-9949-5)
- Grosz B., Gelybó Gy., Churkina G., Haszpra L., Hidy D., Horváth L., Kern A., Kljun N., Machon A., Pásztor L., Barcza Z. (2010) Modeling of biosphere-atmosphere exchange of greenhouse gases: Arable lands. In: Haszpra L. (Ed.) Atmospheric Greenhouse Gases: The Hungarian Perspective. Dordrecht; London; New York: Springer-Science, 2011. pp. 263-294. (ISBN:978-90-481-9949-5)
- Farkas Cs., Alberti G., Balogh J., Barcza Z., Birkás M., Czóbel Sz., Davis K. J., Führer E., Gelybó Gy., Grosz B., Kljun N., Koós S., Machon A., Marjanović H., Nagy Z., Peresotti A., Pintér K., Tóth E., Horváth L. (2010) Measurements and estimations of biosphere-atmosphere exchange of greenhouse gases: Methodologies. In: Haszpra L. (Ed.) Atmospheric Greenhouse Gases: The Hungarian Perspective. Dordrecht; London; New York: Springer-Science, 2011. pp. 65-90. (ISBN:978-90-481-9949-5)
- Barcza Z., Bondeau A., Churkina G., Ciais P., Czóbel Sz., Gelybó Gy., Grosz B., Haszpra L., Hidy D., Horváth L., Machon A., Pásztor L., Somogyi Z., Van Oost K. (2010) Modeling of biosphere-atmosphere exchange of greenhouse gases: Model Based Biospheric Greenhouse Gas Balance of Hungary In: Haszpra L. (Ed.) Atmospheric Greenhouse Gases: The Hungarian Perspective. Dordrecht; London; New York: Springer-Science, 2011. pp. 295-330. (ISBN:978-90-481-9949-5)
- Mihon D., Bacu V., Mészáros R., Gelybó Gy., Gorgan D. (2010) Satellite image interpolation and analysis through GreenView application. In: 4th International Conference on Complex, Intelligent and Software Intensive Systems: CISIS-2010 Krakow, Poland, 15-18 February 2010 pp. 514-519. (ISBN:978-076953967-6)

Conferences

- Kljun N., Gelybó Gy., Barcza, Z., Kern, A. (2013): Effect of spatial heterogeneity on the validation of remote sensing based GPP estimations. GEOPHYSICAL RESEARCH ABSTRACTS 15: Paper EGU2013-8056-2
- Tóth E., Gelybó Gy., Farkas Cs., Lagzi I. (2012) Effect of inter-row cultivation on soil CO₂ emission in a peach plantation GEOPHYSICAL RESEARCH ABSTRACTS 14: Paper EGU2012-12768.
- Mihon D., Bacu V., Gorgan D., Mészáros R., Gelybó Gy., Stefanut T. (2010) GreenView and GreenLand Applications Development on SEE-GRID Infrastructure GEOPHYSICAL RESEARCH ABSTRACTS 12: Paper EGU2010-14455.

- Kljun N., Barcza Z., Kern A., Haszpra L., Gelybó Gy. (2009) Spatial Representativeness of Tall Flux Tower GEOPHYSICAL RESEARCH ABSTRACTS 11: Paper EGU2009-14115.
- Kern A., Bartholy J., Pongrácz R., Barcza Z., Fassang Á., Gelybó Gy. (2007) Analysis of NOAA Pathfinder NDVI time series for Central Europe. GEOPHYSICAL RESEARCH ABSTRACTS 9: Paper EGU2007-00953.
- Gelybó Gy., Barcza Z., Kern A., Kljun N., Haszpra L. (2010) Tall tower based evaluation of the MOD17 model over heterogeneous agricultural vegetation In: AMS 29th Conf. on Agricultural and Forest Meteorology/19th Symp. on Boundary Layers and Turbulence/Ninth Symp. on the Urban Environment. Keystone, USA, 1-6 August 2010.
- Gelybó Gy., Barcza Z., Kern A., Alberti G., Di Tommasi P., Elbers J. A., Fischer M. L., Gonzalez-Meler M., Griffis T. J., Kutsch W. L., Magliulo E. V., Matamala R., Moors E. J., Peressotti A., Suyker A. E., Verma S. B. (2010) Multi-site evaluation of the MOD17 model over homogeneous agricultural vegetation. In: AMS 29th Conf. on Agricultural and Forest Meteorology/19th Symp. on Boundary Layers and Turbulence/Ninth Symp. on the Urban Environment. Keystone, USA, 1-6 August 2010.
- Kljun N., Gelybó Gy., Barcza Z., Kern A., Haszpra L. (2010) Attributing Tall Tower Flux Data to Heterogeneous Vegetation 29th Conference on Agricultural and Forest Meteorology/19th Symposium on Boundary Layers and Turbulence/Ninth Symposium on the Urban Environment, Keystone, Colorado, USA, 1-6 August 2010.
- Barcza Z., Gelybó Gy., Kern A., Kljun N., Haszpra L. (2010). Estimation of the carbon balance components of heterogeneous agricultural landscape using tall tower based and remotely sensed data (poster). 17th International TOVS Study Conference (ITSC-17), Monterey, California, USA, 14-21. April 2010.
- Gelybo Gy., Barcza Z., Kern A., Kljun N., Haszpra L. (2009) Estimation of the carbon balance components of heterogeneous agricultural landscape using tall tower based and remotely sensed data. In: 8. International Carbon Dioxide Conference. Jena, Germany, 13-19 September 2009. Paper T4-008.
- Gelybo Gy., Barcza Z., Haszpra L., Kern A., Tomelleri E., Reichstein M., Bartholy J. (2008) MODIS GPP product: evaluation and calibration in Hungary using tall tower based direct flux measurements. In: 8th EMS Annual Meeting/7th European Conference on Applied Climatology (ECAC) Abstracts Vol 7. Amsterdam, The Netherlands, 29 September-3 October 2008.

Other selected publications

- Bartholy J., Pongrácz R., Gelybó Gy. (2009): Climate signals of the North Atlantic Oscillation detected in the Carpathian Basin. *APPLIED ECOLOGY AND ENVIRONMENTAL RESEARCH*, 7 (3), 229-240.
- Pongrácz R., Bartholy J., Szabó, P., Gelybó Gy. (2009): A comparison of the observed trends and simulated changes in extreme climate indices in the Carpathian Basin by the end of this century. *INTERNATIONAL JOURNAL OF GLOBAL WARMING*, 1 (1/2/3), 336-355. doi: 10.1504/IJGW.2009.027097
- Bartholy J., Pongrácz R., Gelybó Gy., Szabó, P. (2008): Analysis of expected climate change in the Carpathian Basin using the PRUDENCE results. *IDŐJÁRÁS*, 112 (3-4), 249-264.
- Bartholy J., Pongrácz R., Gelybó Gy. (2007): Regional climate change expected in Hungary for 2071-2100. *APPLIED ECOLOGY AND ENVIRONMENTAL RESEARCH*, 5 (1), 1-17.